

Extra exercises in STK4080 2019

Exercise E4.1

- a) Simulate 100 observations of the triple (t, δ, x) by using the commands:

```
x=rgamma(100,2)
T=sqrt(rexp(100)*2*exp(-x))
C=rexp(100,.5)
t=pmin(C,T)
delta=1*(T<C)
```

Write down the density of the variables x and C .

The triple gives a data set of 100 observations of (\tilde{T}_i, D_i, x_i) , according to the standard notation,

Use the code to find an expression for the hazard rate function of the underlying lifetimes T_i . [Answer: $\alpha(t|x_i) = te^{\beta x_i}$]

- b) Fit a Cox-model to the data using, e.g., the command

```
cfit=coxph(Surv(t,delta)^~x) (*)
```

Then plot the martingale residuals with a corresponding lowess smooth (you may follow the setup of Slides 12). Give a comment to the plot.

- c) Now let x and C have the same distributions as before, but simulate new T by

```
T=sqrt(rexp(100)*2*exp(-log(x)))
```

Write down the hazard rate of the new T and put it on the form of Cox-regression with a transformed covariate. What is the transformation of x ?

- d) Then fit a Cox-model using (*) with the new data, thus still assuming the hazard ratio to be $e^{\beta x}$.

Plot the new martingale residuals and the lowess smooth. Comment on the fit.

- e) Finally, try to find the correct form of a transformation of x in the Cox model, i.e., try to find an $f(x)$ such that the hazard ratio is $e^{f(x)}$.

Since there is only one covariate, you should start by fitting an empty model and then look at its martingale residuals.

```
cfit.nox=coxph(Surv(t,delta)^~1)
martres.nox = cfit.nox$residuals
```

Then make a lowess smooth and comment!

Exercise E4.2

In this exercise we will study data simulated from a regression model where the “ β ” depends on time. Let there be a single covariate $x > 0$, drawn from the same distribution as in the previous exercise, and let C now be exponential with hazard rate 0.3. Assume that the true hazard rate is

$$\alpha(t|x) = e^{\beta \log(t)x} = t^{\beta x}$$

- a) Is the given model a proportional hazards model?

Show that for given value of $x > 0$, T has survival function

$$S(t|x) = \exp \left\{ -\frac{t^{\beta x+1}}{\beta x + 1} \right\}$$

and hence is Weibull-distributed with shape parameter $\beta x + 1$ and scale parameter $(\beta x + 1)^{1/(\beta x + 1)}$ (with the parameterization used by R)

- b) The following code will simulate $n = 200$ triples (t, delta, x) with the given distribution for t given x :

```
n=200
x=rgamma(n, 2)
beta=.2
y=beta*x+1
T=rweibull(n,y,y^(1/y))
C=rexp(n,.3)
t=pmin(C,T)
delta=1*(T<C)
```

- c) Fit an ordinary Cox-model with hazard ratio $e^{\beta x}$ to the data and comment on the results. You may also look at martingale residuals.

Then do the test of proportional hazards by using the `cox.zph` function, and draw a scaled Schoenfeld residual plot. Use for example

```
cox.zph(cfit, transform='log')
plot(cox.zph(cfit, transform='log'))
```

(see Slides 12). Comment on the result. (Note that the resulting p -value will vary if you simulate new sets of 200 observations.)

- d) Try other choices for “transform” in the `cox.zph` function. Here is from the R-documentation: **Transform** is a character string specifying how the survival times should be transformed before the test is performed. Possible values are ”km”, ”rank”, ”identity” or a function of one argument.