

ON THE CONSTRUCTION OF THE FORWARD AND BACKWARD SOLUTION SPACE OF A DISCRETE TIME AR-REPRESENTATION

N. P. Karampetakis*

* *Department of Mathematics, Aristotle University of
 Thessaloniki, Thessaloniki 54006, Greece, karampet@auth.gr*

Abstract: The main purpose of this work is to construct the forward and backward solution space of a nonregular discrete time AR-representation i.e. $A(\sigma)\xi(k) = 0$, in a closed interval $[0, N]$ where $A(\sigma)$ is a polynomial matrix and σ is the forward shift operator. The construction of the behavior is based on the structural invariants of the polynomial matrix which describe the AR-representation, i.e. the finite and infinite elementary divisors and the right minimal indices of $A(\sigma)$.

Keywords: Autotegressive models, behaviour, boundary conditions, discrete time systems, indices

Consider a system of linear homogeneous difference and algebraic equations described in matrix form by :

$$A(\sigma)\xi_k = 0 \quad (1)$$

where σ denotes the forward shift operator i.e. $\sigma\xi_k = \xi_{k+1}$, $A(\sigma) = A_0 + A_1\sigma + \dots + A_q\sigma^q \in R[\sigma]^{p \times m}$ with $\text{rank}_{R(\sigma)} A(\sigma) = r \leq \min(p, m)$ and $\xi_k : [0, N] \rightarrow R^m$. We call the set of equations (1) an AR-Representation of B_D (behavior) where B_D is defined as :

$$B_D := \left\{ \begin{array}{l} \xi_k : [0, N] \rightarrow R^m \\ (1) \text{ is satisfied } \forall k \in [0, N] \end{array} \right\}$$

In case where $A(\sigma)$ is regular the solution vector space of (1) consists of rq i.e. the total number of finite and infinite elementary divisors (order accounted for), linearly independent forward and backward solutions (Antoniou *et al.*, 1998). In case now where $A(\sigma)$ is nonregular, the space B_D contains a number of linear independent forward and backward solutions that depends on N and are due to the right null space of $A(\sigma)$ (Karampetakis, 2001). If we now correspond all

the forward and backward solutions which are due to a specific boundary value (initial-final condition) to an element $[\xi_k]$ then the behaviour space B_D is divided into equivalence classes and a new space is created, named \hat{B}_D , whose dimension is proved to be the total number of the finite elementary divisors (n), infinite elementary divisors (μ), plus two times the right minimal indices (ϵ) of $A(\sigma)$ (order accounted for) i.e. $n + \mu + 2\epsilon$ (Karampetakis, 2001). In this paper we give a specific construction of the solution space \hat{B}_D by finding $n + \epsilon$ linearly independent forward solutions representatives due to the finite elementary divisors and the right minimal indices of $A(\sigma)$ and $\mu + \epsilon$ linearly independent backward solution representatives due to the infinite elementary divisors and the right minimal indices of $A(\sigma)$. Finally, the construction of the basis that produces the space \hat{B}_D helps us to construct additionally the space B_D .

1. PRELIMINARY RESULTS

Applying the Z transform of $\xi(k)$ given by (Freeman, 1965) i.e. $\xi(z) \stackrel{def}{=} Z[\xi(k)] = \sum_{k=0}^N \xi(k) z^{-k}$ to (1) we get :

$$\begin{aligned}
& A(z)\xi(z) = [z^q I_p \cdots z I_p I_p] \times \\
& \times \begin{bmatrix} A_q & 0 & \cdots & 0 \\ A_{q-1} & A_q & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ A_1 & A_2 & \cdots & A_q \end{bmatrix} \begin{bmatrix} \xi(0) \\ \xi(1) \\ \vdots \\ \xi(q-1) \end{bmatrix} + \\
& + [z^{-N} I_p \cdots z^{-N+q-2} I_p z^{-N+q-1} I_p] \times \\
& \times \begin{bmatrix} A_0 & 0 & \cdots & 0 \\ A_1 & A_0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ A_{q-1} & A_{q-2} & \cdots & A_0 \end{bmatrix} \begin{bmatrix} \xi(N) \\ \xi(N-1) \\ \vdots \\ \xi(N-q+1) \end{bmatrix} \\
& =: (z^q I_p) \mathfrak{R}_q \tilde{\xi}(0) + (z^{-N} I_p) \mathfrak{R}_0 \tilde{\xi}(N) \quad (2)
\end{aligned}$$

Let also a right minimal polynomial basis¹ consisted by the following polynomial vectors

$$\begin{aligned}
\tilde{u}_i(\sigma) &:= \tilde{u}_{i,0} + \tilde{u}_{i,1}\sigma + \cdots + \tilde{u}_{i,\varepsilon_i}\sigma^{\varepsilon_i} \\
i &= r+1, r+2, \dots, m
\end{aligned}$$

Then the rational vectors of the form

$$\begin{aligned}
\hat{u}_i(\sigma) &:= \tilde{u}_{i,0} \frac{1}{\sigma^{\varepsilon_i}} + \tilde{u}_{i,1} \frac{1}{\sigma^{\varepsilon_i-1}} + \cdots + \tilde{u}_{i,\varepsilon_i} \quad (3) \\
i &= r+1, r+2, \dots, m
\end{aligned}$$

still constitutes a right minimal proper basis of $A(\sigma)$. Let now

$$X = \left\{ \begin{array}{l} \xi(k) : \tilde{\xi}(0) \in Ker[\mathfrak{R}_q] \\ \text{and } \tilde{\xi}(N) \in Ker[\mathfrak{R}_0] \\ \xi(k) = \sum_{i=0}^k \hat{u}_{r+1}(i) z_1(k-i) + \\ + \cdots + \sum_{i=0}^k \hat{u}_m(i) z_{m-r}(k-i) \end{array} \right\}$$

where $z_i(k)$, $i = 1, 2, \dots, m-r$ are arbitrary discrete time functions and $\hat{u}_i(k) = Z^{-1}[\hat{u}_i(z)]$, $i = r+1, \dots, m$, be the solution space of (1) (Karampetakis, 2001) which comes under the above specific initial-final conditions.

Define now the following relation R between the solutions of B_D

$$\begin{aligned}
R(\xi_1(k), \xi_2(k)) &:= \quad (4) \\
&= \left\{ (\xi_1(k), \xi_2(k)) : \xi_1(k) - \xi_2(k) \in X \right. \\
&\quad \left. \text{where } \xi_1(k), \xi_2(k) \in B_D \right\}
\end{aligned}$$

Then the relation (4) defines an equivalence relation (Karampetakis, 2001). We call an equivalence class of the element $\xi(k) \in B_D$, and we denote

this with $[\xi(k)]_R$, the set of all the elements of B_D which are equivalent to $\xi(k)$ or equivalently

$$[\xi(k)]_R := \xi(k) \oplus X \quad (5)$$

We can see that any *equivalence class* of an element $\xi(k)$ gives all the solutions of (1) under some specific initial-final conditions. In the case where $A(\sigma)$ has no right kernel then every equivalence class is composed either by a unique element or no element (due to left null space) contrary to the nonregular case where to each equivalence class corresponds an arbitrary number of elements of B_D . We conclude therefore, that B_D is divided into equivalence classes which are defined by (5). Define now the following *sum* and *product* between equivalence classes of the form (5) :

$$\begin{aligned}
[\xi_1(k)]_R + [\xi_2(k)]_R &:= [\xi_1(k) + \xi_2(k)]_R \\
\lambda [\xi(k)]_R &= [\lambda \xi(k)]_R \quad \lambda \in R
\end{aligned}$$

Theorem 1. (Karampetakis, 2001) The space which is spanned by the equivalence classes defined in (5) is a vector space $\hat{B}_D := B_D/R$ and has dimension

$$f := \dim \hat{B}_D = n + \mu + 2\varepsilon$$

In the following section we shall try to determine a basis for the vector space \hat{B}_D in terms of the structural invariants of the polynomial matrix $A(s)$ and thus to find out the solution space B_D .

2. CONSTRUCTION OF THE BEHAVIOR OF A DISCRETE TIME AR-REPRESENTATION

In this section we shall try to define n linearly independent forward solutions of (1) due to the finite elementary divisors of $A(s)$, μ linearly independent backward solutions of (1) due to the infinite elementary divisors of $A(s)$, ε linearly independent forward solutions of (1) due to the right minimal indices of $A(s)$ and ε linearly independent backward solutions of (1) due to right minimal indices of $A(s)$.

2.1 Finite elementary divisors and solutions of discrete time AR-representations

Let us assume that $A(\sigma)$ has k distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_k$ where for simplicity of notation we assume that $\lambda_i \in R$, $i \in k$ and let

$$\begin{aligned}
U_L(\sigma) A(\sigma) U_R(\sigma) &= \quad (6) \\
&= \text{blockdiag}[I_{z-1}, f_z(\sigma), f_{z+1}(\sigma), \dots, f_r(\sigma), 0_{p-r, m-r}]
\end{aligned}$$

¹ For the simplicity of the proofs of the main Theorems we give a specific construction of the minimal polynomial basis in (7), although in practice other constructions may also be used i.e. (Beelen, 1987).

$1 \leq z \leq r$ be the Smith form of $A(\sigma)$ (in C) where $f_i(\sigma) \in R[\sigma]$ are the invariant polynomials of $A(\sigma)$ and $f_i(\sigma)/f_{i+1}(\sigma)$ $i = z, z+1, \dots, r-1$. Assume that the partial multiplicities of the eigenvalues $\lambda_i \in R, i \in k$ are $0 \leq \sigma_{i,z} \leq \sigma_{i,z+1} \leq \dots \leq \sigma_{i,r}$. Let $u_j(\sigma) \in R[\sigma]^{m \times 1}, j \in R$ be the columns of $U_R(\sigma)$ and $u_j^{(q)}(\sigma) := (d^q/ds^q) u_j(\sigma), q = 0, 1, \dots, \sigma_{i,j} - 1$. Let also

$$x_{j,q}^i := \frac{1}{q!} u_j^{(q)}(\lambda_i) \quad i \in k \text{ and } j = z, z+1, \dots, r$$

Define the vector valued functions

if $\lambda_i \neq 0$

$$\xi_{j,q}^i(k) := \lambda_i^k x_{j,q}^i + \dots + \binom{k}{q} \lambda_i^{k-q} x_{j,0}^i$$

$$i \in k; j = z, z+1, \dots, r; q = 0, 1, \dots, \sigma_{i,j} - 1$$

if $\lambda_i = 0$

$$\xi_{j,q}^i(k) := \Delta(k) x_{j,q}^i + \dots + \Delta(k-q) x_{j,0}^i$$

$$i \in k; j = z, z+1, \dots, r; q = 0, 1, \dots, \sigma_{i,j} - 1$$

Let

$$\Psi_{i,j}(k) := [\xi_{j,0}^i(k) \ \xi_{j,1}^i(k) \ \dots \ \xi_{j,\sigma_{i,j}-1}^i(k)]$$

$$C_{i,j} := [x_{j,0}^i \ x_{j,1}^i \ \dots \ x_{j,\sigma_{i,j}-2}^i \ x_{j,\sigma_{i,j}-1}^i]$$

$$J_{i,j} := \begin{bmatrix} \lambda_i & 1 & \dots & 0 & 0 \\ 0 & \lambda_i & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & \lambda_i & 1 \\ 0 & 0 & \dots & 0 & \lambda_i \end{bmatrix} \in R^{\sigma_{i,j} \times \sigma_{i,j}}$$

$$i \in k; j = z, z+1, \dots, r$$

and

$$\Psi_i^F(k) := [\Psi_{i,z}(k) \ \Psi_{i,z+1}(k) \ \dots \ \Psi_{i,r}(k)]$$

$$C_i^F := [C_{i,z}(k) \ C_{i,z+1}(k) \ \dots \ C_{i,r}(k)]$$

$$J_i^F := \text{blockdiag} [J_{i,z}(k) \ J_{i,z+1}(k) \ \dots \ J_{i,r}(k)]$$

where $m_i := \sigma_{i,z} + \sigma_{i,z+1} + \dots + \sigma_{i,r}$ $i \in k$. Finally let

$$\Psi_F^D(k) := [\Psi_1^F(k) \ \Psi_2^F(k) \ \dots \ \Psi_r^F(k)]$$

$$C_F^D := [C_1^F(k) \ C_2^F(k) \ \dots \ C_r^F(k)]$$

$$J_F^D := \text{blockdiag} [J_1^F(k) \ J_2^F(k) \ \dots \ J_r^F(k)]$$

where $n := \deg \left[\prod_{j=z}^r f_j(\sigma) \right] = \deg \left| S_{R(\sigma)}^C \right|$. Then we have the following

Theorem 2. The columns of the following matrix

$$\Psi_F^D(k) := [\Psi_1^F(k) \ \Psi_2^F(k) \ \dots \ \Psi_r^F(k)] = C_F^D (J_F^D)^k$$

constitute a solution subspace B_F^D of $B_D - X$ i.e. $B_F^D \subseteq B_D - X$ with dimension

$$\dim B_F^D = n := \left\{ \begin{array}{l} \text{total sum of the degrees of the} \\ \text{finite elementary divisors of } A(\sigma) \end{array} \right\}$$

Proof. It is easily seen (see also (Gogberg *et al.*, 1982)) that the pair (C_F^D, J_F^D) constitute a finite spectral pair of $A(\sigma)$ which satisfy the following :

$$\sum_{k=0}^q A_k C_F^D (J_F^D)^k = 0; \text{rank col} \left(C_F^D (J_F^D)^k \right)_{k=0}^{n-1} = n$$

and, therefore, the columns of the matrix $\Psi_F^D(k)$ satisfy the equation (1). In order to prove that $B_F^D \subseteq B_D - X$ we have to show that the columns of the matrix $\Psi_F^D(k)$ do not belong to X or otherwise

$$\left[C_F^T (C_F J_F)^T \ \dots \ (C_F J_F^{q-1})^T \right]^T \notin \text{Ker} [R_q]$$

$$\left[(C_F J_F^N)^T (C_F J_F^N)^{N-1} \ \dots \ (C_F J_F^N)^{N-q} \right] \notin \text{Ker} [R_0]$$

The proof is not presented here in order to avoid including a lot of technicalities. However, it is based a) on the specific selection of the right minimal basis selected in the sequel and b) on the linearly independence of the columns of the transforming matrix $U_R(s)$ defined above. The proof follows similar lines to the ones for the continuous time case presented in (Karampetakis, 1993) (Chapter 6).

Any other finite spectral pair will also define an isomorphic space to B_F^D . However our intention is twofold : a) to give the reader a method for the construction of the finite spectral pair and b) to simplify some of the proofs with the specific form of this spectral pair.

2.2 Infinite elementary divisors and solutions of discrete time AR-representations

Define the "dual" polynomial matrix $\tilde{A}(w)$ of $A(\sigma)$ as

$$\tilde{A}(w) := A_0 w^q + A_1 w^{q-1} + \dots + A_q \in R[w]^{p \times m}$$

Let $\tilde{U}_L(w) \in R(w)^{p \times p}, \tilde{U}_R(w) \in R(w)^{m \times m}$ be rational matrices having no poles and zeros at $w = 0$ and such that

$$\begin{aligned} \tilde{U}_L(w)\tilde{A}(w)\tilde{U}_R(w) &= S_{\tilde{A}(w)}^0(w) = \\ &= \text{blockdiag}[I_{d-1}, z^{\mu_d}, \dots, z^{\mu_r}, 0_{p-r, m-r}] \end{aligned}$$

where $S_{\tilde{A}(w)}^0(w)$ is the Smith form of $\tilde{A}(w)$ at $w = 0$. Let now $\tilde{U}_R(w) = [\tilde{u}_1(w) \ \tilde{u}_2(w) \ \cdots \ \tilde{u}_m(w)]$ where $\tilde{u}_j(w) \in R(w)^{m \times 1}$ and $\tilde{u}_j^{(i)}(w), \tilde{A}^{(i)}(w)$ be the q th derivatives of $\tilde{u}_j(w)$ and $\tilde{A}(w)$ with respect to w for $i = 0, 1, \dots, \mu_j - 1$ and $j = d, d+1, \dots, r$ where μ_j are the multiplicities of the zeros of $\tilde{A}(w)$ at $w = 0$ or equivalently the degrees of the infinite elementary divisors. Define

$$x_{j,i} := \frac{1}{i!} \tilde{u}_j^{(i)}(0)$$

for $i := 0, 1, \dots, \mu_j - 1$ and $j = d, d+1, \dots, r$. Then for final conditions

$$\begin{bmatrix} \xi(N) \\ \xi(N-1) \\ \vdots \\ \xi(N-q+1) \end{bmatrix} = \begin{bmatrix} x_{j,i} \\ \vdots \\ x_{j,0} \\ 0_{(q-i-1),1} \end{bmatrix}$$

we obtain respectively the linearly independent backward solutions

$$\xi_{j,i}^B(k) := x_{j,i} \delta(N-k) + \cdots + x_{j,0} \delta(N-(k-i))$$

Define the vector valued functions

$$\begin{aligned} \xi_{j,i}^B(k) &:= x_{j,i} \delta(N-k) + \cdots + x_{j,0} \delta(N-(k-i)) \\ i &:= 0, 1, \dots, \mu_j - 1 \text{ and } j = k, k+1, \dots, r \end{aligned}$$

Let

$$\begin{aligned} \Psi_j^B(k) &:= \begin{bmatrix} \xi_{j,0}^B(k) & \xi_{j,1}^B(k) & \cdots & \xi_{j,\mu_j-1}^B(k) \end{bmatrix} \\ C_j^B &:= \begin{bmatrix} x_{j,0} & x_{j,1} & \cdots & x_{j,\mu_j-1} \end{bmatrix} \\ J_j^B &:= \begin{bmatrix} 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 0 \end{bmatrix} \in R^{\mu_j \times \mu_j} \end{aligned}$$

where $j = d, d+1, \dots, r$ and

$$\begin{aligned} \Psi_B^D(k) &:= [\Psi_k^B(k) \ \Psi_{k+1}^B(k) \ \cdots \ \Psi_r^B(k)] \\ C_B^D &:= [C_k^B(k) \ C_{k+1}^B(k) \ \cdots \ C_r^B(k)] \\ J_B^D &:= \text{blockdiag} [J_k^B(k) \ J_{k+1}^B(k) \ \cdots \ J_r^B(k)] \end{aligned}$$

where $\mu := \sum_{j=d}^r \mu_j$. Then we have the following

Theorem 3. The columns of the following matrix

$$\Psi_B^D(k) := [\Psi_k(k) \ \Psi_{k+1}(k) \ \cdots \ \Psi_r(k)] = C_B^D (J_B^D)^{N-k}$$

constitute a solution subspace B_B^D of $B_D - X$ i.e. $B_B^D \subseteq B_D - X$ with dimension

$$\dim B_B^D = \mu := \left\{ \begin{array}{l} \text{total sum of the degrees of the} \\ \text{infinite elementary divisors} \end{array} \right\}$$

Proof. It is easily seen that the pair (C_B^D, J_B^D) constitutes an infinite spectral pair of $A(\sigma)$ which satisfies the following :

$$\sum_{k=0}^q A_k C_B^D (J_B^D)^{N-k} = 0 ; \text{rank col} \left(C_B^D (J_B^D)^{N-k} \right)_{k=0}^{\mu-1} = \mu$$

and, therefore, the columns of the matrix $\Psi_B^D(k)$ satisfy the equation (1). Similar comments to the ones in the proof of Theorem 2, also apply here.

Any other infinite spectral pair which corresponds to infinite elementary divisors will also define an isomorphic space to B_B^D .

2.3 Right minimal indices and solutions of discrete time AR-representations

$A(\sigma) \in R[\sigma]^{p \times m}$ according to our assumption has rank $r \leq \min\{p, m\}$ and therefore the dimension of the right null space of $A(\sigma)$ is equal to $m - r$. Consider a minimal polynomial bases² of the right null space of $A(\sigma)$, let

$$[\bar{u}_{r+1}(\sigma) \ \bar{u}_{r+2}(\sigma) \ \cdots \ \bar{u}_m(\sigma)] \quad (7)$$

The greatest degrees of the columns $\bar{u}_i(\sigma), i = r+1, r+2, \dots, m$ let $\{\varepsilon_{r+1}, \varepsilon_{r+2}, \dots, \varepsilon_m\}$ are called right minimal indices of $A(\sigma)$. Let also

$$x_{j,i} := \frac{1}{i!} \bar{u}_j^{(i)}(0)$$

$$i = 0, 1, \dots, \varepsilon_i - 1 \text{ and } j = r+1, r+2, \dots, m$$

Define the vector valued functions

$$\begin{aligned} \xi_{j,i}^F(k) &:= \delta(k) x_{j,i} + \cdots + \delta(k-i) x_{j,0} \\ i &:= 0, 1, \dots, \varepsilon_j - 1 \text{ and } j = r+1, r+2, \dots, m \end{aligned}$$

Let

² The last $m - r$ columns of the transforming matrix $U_R(\sigma)$ defined in subsection 2.1, constitute a basis of the right kernel of $A(\sigma)$. Under certain unimodular transformations i.e. column reduceness, we may do the above basis minimal.

$$\begin{aligned}\Psi_j^F(k) &:= [\xi_{j,0}^F(k) \ \xi_{j,1}^F(k) \ \cdots \ \xi_{j,\varepsilon_j-1}^F(k)] \\ C_j^F &:= [x_{j,0} \ x_{j,1} \ \cdots \ x_{j,\varepsilon_j-1}] \\ J_j^F &:= \begin{bmatrix} 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 0 \end{bmatrix} \in R^{\varepsilon_j \times \varepsilon_j}\end{aligned}$$

where $j = r+1, r+2, \dots, m$ and

$$\begin{aligned}\Psi_F^\varepsilon(k) &:= [\Psi_{r+1}^F(k) \ \Psi_{r+2}^F(k) \ \cdots \ \Psi_m^F(k)] \\ C_F^\varepsilon &:= [C_{r+1}^F(k) \ C_{r+2}^F(k) \ \cdots \ C_m^F(k)] \\ J_F^\varepsilon &:= \text{blockdiag} [J_{r+1}^F(k) \ J_{r+2}^F(k) \ \cdots \ J_m^F(k)]\end{aligned}$$

where $\varepsilon := \sum_{j=r+1}^m \varepsilon_j$. Then we have the following

Theorem 4. The columns of the following matrix

$$\begin{aligned}\Psi_F^\varepsilon(k) &:= C_F^\varepsilon (J_F^\varepsilon)^k = \\ &= [\Psi_{r+1}^F(k) \ \Psi_{r+2}^F(k) \ \cdots \ \Psi_m^F(k)]\end{aligned}$$

constitute a solution subspace B_F^ε of B_D i.e. $B_F^\varepsilon \subseteq B_D - X$ with dimension

$\dim B_F^\varepsilon = \varepsilon :=$ total sum of the right minimal indices

Proof. Similar to the proof of Theorem 2.

Consider the dual minimal polynomial base of (7). It is easily seen that constitutes a minimal bases of the right null space of the dual polynomial matrix $\tilde{A}(\sigma)$ of $A(\sigma)$, let

$$[\tilde{u}_{r+1}(\sigma) \ \tilde{u}_{r+2}(\sigma) \ \cdots \ \tilde{u}_m(\sigma)]$$

The greatest degrees of the columns $\tilde{u}_i(\sigma)$, $i = r+1, r+2, \dots, m$ are the same as the right minimal indices of $A(\sigma)$ i.e. $\{\varepsilon_{r+1}, \varepsilon_{r+2}, \dots, \varepsilon_m\}$.³ Let also

$$x_{j,i} := \frac{1}{i!} \tilde{u}_j^{(i)}(0) \quad i = 0, 1, \dots, \varepsilon_j - 1 \text{ and } j = r+1, r+2, \dots, m$$

Define the vector valued functions

$$\begin{aligned}\xi_{j,i}^B(k) &:= \delta(N-k) x_{j,i} + \cdots + \delta(N-(k-i)) x_{j,0} \\ i &= 0, 1, \dots, \varepsilon_j - 1 \text{ and } j = r+1, r+2, \dots, m\end{aligned}$$

Let

³ The vectors $u_i(s)$ $i = r+1, \dots, m$ are linearly independent and thus its values at $s = 0$ i.e. $u_i(0)$ $i = r+1, \dots, m$, are also linearly independent. Therefore the leading coefficient matrix of the vectors $\tilde{u}_i(s)$ of the dual polynomial basis are not zero and have the same degrees as the ones of the right minimal polynomial basis of $A(\sigma)$.

$$\begin{aligned}\Psi_j^B(k) &:= [\xi_{j,0}^B(k) \ \xi_{j,1}^B(k) \ \cdots \ \xi_{j,\varepsilon_j-1}^B(k)] \\ C_j^B &:= [x_{j,0} \ x_{j,1} \ \cdots \ x_{j,\varepsilon_j-1}] \\ J_j^B &:= \begin{bmatrix} 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 0 \end{bmatrix} \in R^{\varepsilon_j \times \varepsilon_j}\end{aligned}$$

where $j = r+1, r+2, \dots, m$ and

$$\begin{aligned}\Psi_B^\varepsilon(k) &:= [\Psi_{r+1}^B(k) \ \Psi_{r+2}^B(k) \ \cdots \ \Psi_m^B(k)] \\ C_B^\varepsilon &:= [C_{r+1}^B(k) \ C_{r+2}^B(k) \ \cdots \ C_m^B(k)] \\ J_B^\varepsilon &:= \text{blockdiag} [J_{r+1}^B(k) \ J_{r+2}^B(k) \ \cdots \ J_m^B(k)]\end{aligned}$$

Then we have the following

Theorem 5. The columns of the following matrix

$$\begin{aligned}\Psi_B^\varepsilon(k) &:= C_B^\varepsilon (J_B^\varepsilon)^{N-k} = \\ &= [\Psi_{r+1}^B(k) \ \Psi_{r+2}^B(k) \ \cdots \ \Psi_m^B(k)]\end{aligned}$$

constitute a solution subspace B_B^ε of B i.e. $B_B^\varepsilon \subseteq B_D - X$ with dimension

$\dim B_B^\varepsilon = \varepsilon :=$ total sum of the right minimal indices

Proof. Similar to the proof of Theorem 3.

Therefore, we conclude that the right minimal indices give rise to ε linearly independent forward solutions and ε linearly independent backward solutions which do not belong to X .

2.4 Construction of the whole solution space

Define the following set of solutions

$$\begin{aligned}\tilde{B}_F^D &= B_F^D \oplus X \quad ; \quad \tilde{B}_B^D = B_B^D \oplus X \\ \tilde{B}_F^\varepsilon &= B_F^\varepsilon \oplus X \quad ; \quad \tilde{B}_B^\varepsilon = B_B^\varepsilon \oplus X\end{aligned}$$

where $B_F^D(k)$, $B_B^D(k)$, $B_F^\varepsilon(k)$ and $B_B^\varepsilon(k)$ are the solution vector spaces which are due to the finite and infinite elementary divisors and the right kernel of $A(\sigma)$. Let also the following spaces

$$\begin{aligned}\hat{B}_F^D &:= \{[\xi(k)]_R : \xi(k) \in \tilde{B}_F^D\} = \tilde{B}_F^D/R \\ \hat{B}_F^\varepsilon &:= \{[\xi(k)]_R : \xi(k) \in \tilde{B}_F^\varepsilon\} = \tilde{B}_F^\varepsilon/R \\ \hat{B}_B^D &:= \{[\xi(k)]_R : \xi(k) \in \tilde{B}_B^D\} = \tilde{B}_B^D/R \\ \hat{B}_B^\varepsilon &:= \{[\xi(k)]_R : \xi(k) \in \tilde{B}_B^\varepsilon\} = \tilde{B}_B^\varepsilon/R\end{aligned}$$

